

New Hadrons Formed by the Fourth SM Family and Iso-singlet Quarks

Hakan CİFTÇİ^a and Saleh SULTANSOY^{a,b}

^a *Department of Physics, Faculty of Arts and Science, Gazi University,
06500 Teknikokullar, Ankara, TURKEY*

^b *Institute of physics, Academy of sciences, H. Cavid Ave 33,
Baku, AZERBAIJAN*

February 1, 2008

Abstract

The properties of new heavy mesons containing new heavy quarks have been investigated. As an example, the fourth SM family quarks and weak iso-singlet quarks predicted by E_6 GUT are considered. Production of these hadrons at TeV energy lepton colliders have been analyzed.

1 Introduction

It is known that t quark does not form meson and baryon states because of the high value of m_t and full strength of tbW vertex. On the other hand, there are strong reasons that the fourth Standard Model (SM) family should exist [1,2]. We expect that the masses of the fourth family quarks lie between 300 and 700 GeV with preferable value $m_4 = 4g_W\eta = 8m_W \cong 640$ GeV [3]. New heavy quarks are also predicted by various extensions of SM and the most wide-known examples are weak iso-singlet quarks predicted by E_6 GUT [4], which is favored by super string theory [5]. In spite of the fact that the masses of new quarks are larger than m_t , they can form hadrons because of the smallness of mixing between new heavy and ordinary quarks. Indeed, according to the parametrization of mass matrices given in [6], mixing between the fourth and third family quarks is predicted to be $|V_{qU_4}| \approx 10^{-3}$. Similar situation is expected for iso-singlet quarks. The condition for forming new hadrons states is [7]

$$m_Q \leq (100\text{GeV}) |V_{Qq}|^{-2/3} . \quad (1)$$

New quarks will be copiously produced at the LHC. The observation of the fourth SM family quarks in ATLAS has been considered in [8, 9]. In this paper, we discuss the properties of new heavy mesons formed by new quarks. In section 2, the properties and the systematics of new hadrons have been investigated. Production of these hadrons at TeV energy lepton colliders have been considered in section 3. Finally, we give some concluding remarks.

2 New Heavy Mesons

According to the Eq. (1), fourth SM family and E_6 isosinglet quarks have formed hadron states if their mixing with known quarks is sufficiently small. In the case of fourth family quarks, parametrization given in [6] satisfies this condition, whereas new hadrons are not formed in the case of parametrization given in [1]. Concerning E_6 isosinglet quarks, we do not have similar parametrization (in this case one deals with 6×6 mass matrix) and one can make only qualitative estimations. For example, if the lightest isosinglet quark has the mass $m_D \cong 0.5$ TeV, new heavy hadrons are formed for $|V_{qD}| < 0.09$.

In order to calculate the masses of new mesons we use the logarithmic potential

$$V(r) = A \log(r) + V_0 , \quad (2)$$

where r is the radial distance between quarks. For two particles system, Schrodinger equation has the form ($\hbar = c = 1$)

$$\left[\frac{p^2}{2\mu} + V(r) \right] \Psi(r) = E \Psi(r) , \quad (3)$$

where $\mu = m_1 m_2 / (m_1 + m_2)$. One can find the binding energy of meson for arbitrary n, l states as

$$E_{nl} \cong V_0 + A \eta_{nl} , \quad (4)$$

where η_{nl} is solution of the equation

$$\left[-\frac{d^2}{dr^2} + \log(r) + \frac{l(l+1)}{r^2} \right] g(r) = \eta g(r). \quad (5)$$

Below, instead of numerical calculations, we use simplified analytical procedure (for details see [10]) to solve Eq. (5), which gives the results coinciding with numerical ones with precision of order 10^{-3} . Thus the bound state mass of the $q_i \bar{q}_j$ system is

$$M(q_i \bar{q}_j)_{nl} = m_{q_i} + m_{q_j} + V_0 + A\eta_{nl} \quad (6)$$

The calculations involve the potential parameters of the model (A, V_0) and the quark masses ($m_u = m_d, m_s, m_c, m_b$). The potential parameters and the quark masses are obtained from the experimental masses of $\Psi(1S)$, $\Psi(2S)$, $\Upsilon(1S)$, $\Upsilon(2S)$, $D^0(1S)$ and $s\bar{s}(1S)$ by fitting Eq. (6): (A, V_0) = (0.7328 GeV, -0.8684 GeV), $m_u = m_d = 0.367$ GeV, $m_s = 0.561$ GeV, $m_c = 1.6$ GeV, $m_b = 4.7816$ GeV. In addition to these parameters, the fourth SM family up-quark and E_6 isosinglet quark masses are chosen as 638.6 GeV [6] and 0.5 TeV, respectively. With these parameters, we calculate the masses of bound states of $c\bar{c}$ and $b\bar{b}$, which are given in Table 1. It is seen that our results are in good agreement with experimental data [11] (for comparison see [12, 13]).

In Table 2 we present the masses of quarkonia and mesons formed by the fourth SM family up-quark and E_6 isosinglet quark, respectively. It is seen that the decay of $3S$ quarkonia states into mesons containing u and d quarks is admitted. In the case of $4S$ quarkonia decays into mesons containing s quark is added. In our opinion, the best mechanism for the production of heavy mesons formed by u_4 and D quarks is the resonance formation of $3S$ and $4S$ quarkonia at lepton colliders with subsequent decay into corresponding meson-antimeson states.

3 Production of New Mesons at Future Lepton Colliders

The spin averaged Breit-Wigner cross section for a spin- J resonance produced in the collision of the particles of spin S_1 and S_2 [11] is

$$\sigma_{BW}(E) = \frac{2J+1}{(2S_1+1)(2S_2+1)} \frac{4\pi}{k^2} \frac{B_{in}B_{out}\Gamma_{tot}^2}{4(E-R_R)^2 + \Gamma_{tot}^2} \quad (7)$$

where k is the *c.m.* momentum, E is the *c.m.* energy, and B_{in} and B_{out} are the branching fractions of the resonance into the entrance and exit channels. In our case, at $E = E_R$ Eq. (7) takes the form

$$\sigma^{res} = \frac{12\pi}{M^2} B_{in}B_{out} \quad (8)$$

where M is the mass of corresponding quarkonium.

Since lepton colliders have the certain energy spread, the average cross section at $\sqrt{s} \approx M$ can be estimated from

$$\sigma^{ave} \cong \frac{\Gamma_{tot}}{\Delta E_{coll}} \sigma^{res}. \quad (9)$$

Here, we take into account that $\Gamma_{tot} \ll \Delta E_{coll}$. Indeed, $\Delta E_{coll} = O(1 \text{ GeV})$ for muon collider [14] and $\Delta E_{coll} = O(10 \text{ GeV})$ for CLIC [15]. Concerning Γ_{tot} , we use Coulomb potential in order to estimate partial decay widths to e^+e^- and W^+W^- (which is dominant one among the decays into fundamental SM particles [16, 17]) taking into account that distance between quarks and antiquarks is much less than 1 fm due to large value of new quarks masses. Results are presented in Table 3. Then, decay width of 3S levels into meson-antimeson states is expected to be $O(100 \text{ MeV})$. Multiplying average cross-sections with integrated luminosity, which are 50 fb^{-1} per year for muon collider [14] and 200 fb^{-1} per year for CLIC [15], one can easily obtain numbers of new mesons produced via formation and decays of 3S resonance per working year, namely, ~ 1000 at muon collider and ~ 400 at CLIC for mesons containing u_4 and first family quarks. Corresponding numbers for mesons formed by D and first family quarks are ~ 340 and ~ 140 , respectively.

If these new mesons are sufficiently long-lived (which means very small mixing of heavy quarks with light quarks [18,19]) we will observe corresponding traces in the detector. The decay length is given by $\beta\gamma c\tau$. In our case, $\gamma \cong 1$ and $\beta \cong 6 \cdot 10^{-4}$. For mesons containing u_4 quarks (here we assume the dominance of mixing between fourth and third SM families and take into account that $m_{u_4} \gg m_{W,Z}$)

$$\tau \approx \frac{8x_W}{\alpha_{em} |V_{u_4b}|^2} \times \frac{m_W^2}{m_{u_4}^3} \quad (10)$$

where $x_W = \sin^2 \theta_W \approx 0.21$. Therefore, $l = 6 \cdot 10^{-2} \text{ m} / |V_{u_4b}|^2$ and taking into account present experimental resolution $\sim 100 \mu\text{m}$ we conclude that $|V_{u_4b}|$ should be less than $2.5 \cdot 10^{-9}$. Similar consideration for meson containing D quarks leads to $\sin \varphi = |V_{Du}| < 1.5 \cdot 10^{-9}$ (here we assume the dominance of $D-d$ mixing and taking account flavor changing interactions [5,20]). In our opinion, this scenario is hardly to be realized.

Let us finish this section with some remarks on mesons containing D quark. As mentioned above, in this case, flavor changing neutral current appear at tree level and for $m_D - m_W \approx m_D - m_Z \gg m_Z - m_W$ one has $Br(D \rightarrow u+W) \approx 0.6$ and $Br(D \rightarrow u+Z) \approx 0.4$ [20]. Therefore, we expect $Br(D \rightarrow jet + l^+l^-) \approx 0.012$ and $Br(D \rightarrow jet + \nu\bar{\nu}) \approx 0.072$ for decay modes which differ isosinglet quark from the fourth SM family quarks. For a most spectacular cases, namely one of the mesons decays into jet and l^+l^- pairs ($l = e, \mu$) or jet and $\nu\bar{\nu}$ pairs and the other one decays into three jets we expect ~ 60 events per working year at muon collider and ~ 20 events at CLIC.

4 Conclusion

We show that mesons formed by new heavy quarks can be observed at future lepton colliders due to formation (and corresponding decay) of heavy quarkonia. The numbers of events are not huge (hundreds events comparing to tens thousands $Q\bar{Q}$ pairs which will be produced at LHC [8, 9]). However, pure experimental environment provide an advantage for clarifying the properties of new hadrons.

5 References

1. A. Datta and S. Raychaudhuri, Phys. Rev. D 49 (1994) 4762.
2. A. Celikel, A. K. Ciftci and S. Sultansoy, Phys. Lett. B 342 (1995) 257.
3. S. Sultansoy, arXiv:hep-ph/0004271 (2000).
4. F. Gursey, P. Ramond and P. Sikivie, Phys. Lett. B 60 (1976) 177.
5. J.L. Hewett and T.G. Rizzo, Phys. Reports 183 (1989) 193.
6. S. Atag et al., Phys. Rev. D 54 (1996) 5745.
7. I. Bigi et al., Phys. Lett. B 181 (1986) 157.
8. E. Arik et al., Phys. Rev. D 58 (1998) 117701.
9. ATLAS Detector and Physics Performance TDR, CERN/LHCC/99-15 (1999).
10. H. Ciftci, E. Ataser and H. Kori, in preparation.
11. D.E. Groom et al., Review of Particle Physics, Eur. Phys. J. C 15 (2000).
12. N. Barik, S. N. Jena and D. P. Rath, Phys. Rev. D 41 (1990) 1568.
13. R.N. Faustov, V.O. Galkin, A.V. Tatarintsev and A.S. Vshivtsev, Int. J. Mod. Phys. A 15 (2000) 209.
14. <http://pubweb.bnl.gov/users/bking/www/mucoll/overview.pdf>
15. <http://cern.web.cern.ch/CERN/Divisions/PS/CLIC/Report/Appendix.html>
16. V. Barger et al., Phys. Rev. D 35 (1987) 3366.
17. A. K. Ciftci, R. Ciftci and S. Sultansoy, arXiv:hep-ph/0106222 (2001).
18. P. H. Frampton and P. Q. Hung, Phys. Rev. D 58 (1998) 057704.

19. P. H. Frampton, P. Q. Hung and M. Sher, Phys. Rep. 330 (2000) 263.
20. S. Sultansoy, Turkish J. of Phys. 22 (1998) 575.

Table.1. The masses of the $c\bar{c}$ and $b\bar{b}$ bound states (in MeV).

Level	$c\bar{c}$ calculated	$c\bar{c}$ experiment [11]	$b\bar{b}$ calculated	$b\bar{b}$ experiment [11]
1S	3097	3096.87 ± 0.04	9461	9460.30 ± 0.26
2S	3686	3685.96 ± 0.09	10050	10023.26 ± 0.31
3S	4010	4040 ± 10	10374	10355.2 ± 0.5
4S	4234	4159 ± 20	10598	10580.0 ± 3.5
5S	4405	4415 ± 9	10769	10865 ± 8
1P	3523	3510.51 ± 0.12	9898	9892.7 ± 0.6
2P	3888	-	10272	10268.5 ± 0.4

Table 2. The masses of the bound states formed by the fourth SM family u_4 and isosinglet D quarks (in GeV).

Level	$u_4\bar{u}_4$	$u_4\bar{u}$	$u_4\bar{s}$	$D\bar{D}$	$D\bar{u}$	$D\bar{s}$
1S	1277.10	638.86	639.06	999.90	500.26	500.46
2S	1277.69	639.45	639.65	1000.49	500.85	501.05
3S	1278.01	639.78	639.97	1000.81	501.18	501.37
4S	1278.23	640.00	640.20	1001.03	501.40	501.60

Table 3. Partial decay widths of heavy quarkonia to e^-e^+ (in keV) and W^-W^+ (in MeV). 1 and 2 correspond to $u_4\bar{u}_4$ and $D\bar{D}$, respectively.

Level	Mass(1)	$\Gamma_{e^+e^-}(1)$	$\Gamma_{W^+W^-}(1)$	Mass(2)	$\Gamma_{e^+e^-}(2)$	$\Gamma_{W^+W^-}(2)$
1S	1277.10	35	400	999.90	6.6	142
2S	1277.69	4.4	50	1000.49	0.8	18
3S	1278.01	1.3	15	1000.81	0.24	5
4S	1278.23	0.5	6.2	1001.03	0.1	2.2